Variation of the group 5 grass pollen allergen content of airborne pollen in relation to geographic location and time in season

The HIALINE working group: Jeroen Buters, PhD,^{a,b} Marje Prank, MSc,^c Mikhail Sofiev, PhD,^c Gudrun Pusch, Dipl Ing,^{a,b} Roberto Albertini, PhD,^d Isabella Annesi-Maesano, MD, DSc, PhD,^{e,f} Celia Antunes, PhD,^{g,h} Heidrun Behrendt, MD,^{a,b} Uwe Berger, MBA,ⁱ Rui Brandao, PhD,^g† Sevcan Celenk, PhD,ⁱ Carmen Galan, PhD,^k Łukasz Grewling, PhD,¹ Bogdan Jackowiak, PhD,¹ Roy Kennedy, PhD,^m Auli Rantio-Lehtimäki, PhD,ⁿ Gerald Reese, PhD,^o Ingrida Sauliene, PhD,^p Matt Smith, PhD,ⁱ Michel Thibaudon, Pharm,^q Bernhard Weber, PhD,^o and Lorenzo Cecchi, MD^r Munich and Reinbek, Germany, Helsinki and Turku, Finland, Parma and Florence, Italy, Paris and Brussieu, France, Évora and Coimbra, Portugal, Vienna, Austria, Gorukle-Bursa, Turkey, Cordoba, Spain, Poznan, Poland, Worcester, United Kingdom, and Siauliai, Lithuania

Background: Allergies to grass pollen are the number one cause of outdoor hay fever. The human immune system reacts with symptoms to allergen from pollen.

Objective: We investigated the natural variability in release of the major group 5 allergen from grass pollen across Europe. Methods: Airborne pollen and allergens were simultaneously collected daily with a volumetric spore trap and a high-volume

†Deceased.

This publication arises from HIALINE, a project that has received funding from the European Union in the framework of the Health Programme.

Available online May 6, 2015.

0091-6749/\$36.00

© 2015 American Academy of Allergy, Asthma & Immunology http://dx.doi.org/10.1016/j.jaci.2015.01.049 cascade impactor at 10 sites across Europe for 3 consecutive years. Group 5 allergen levels were determined with a Phl p 5-specific ELISA in 2 fractions of ambient air: particulate matter of greater than 10 µm in diameter and particulate matter greater than 2.5 µm and less than 10 µm in diameter. Mediator release by ambient air was determined in FceRI-humanized basophils. The origin of pollen was modeled and condensed to pollen potency maps. Results: On average, grass pollen released 2.3 pg of Phl p 5 per pollen. Allergen release per pollen (potency) varied substantially, ranging from less than 1 to 9 pg of Phl p 5 per pollen (5% to 95% percentile). The main variation was locally day to day. Average potency maps across Europe varied between years. Mediator release from basophilic granulocytes correlated better with allergen levels per cubic meter ($r^2 = 0.80, P < .001$) than with pollen grains per cubic meter ($r^2 = 0.61, P < .001$). In addition, pollen released different amounts of allergen in the non-pollenbearing fraction of ambient air, depending on humidity. Conclusion: Across Europe, the same amount of pollen released substantially different amounts of group 5 grass pollen allergen. This variation in allergen release is in addition to variations in pollen counts. Molecular aerobiology (ie, determining allergen in ambient air) might be a valuable addition to pollen counting. (J Allergy Clin Immunol 2015;136:87-95.)

Key words: Allergen, grass, Phl p 5, Europe, PM10, PM2.5, exposure, allergy, basophil, modeling, SILAM, HIALINE

Grass pollen is the major cause of allergic sensitizations to outdoor allergens all over Europe¹ and ranks second to house dust mite as a cause of respiratory allergy in most countries.²⁻⁵ The role of grass pollen in respiratory allergy was recently confirmed in a large epidemiologic study from the European Community Respiratory Health Study.⁶

Grass pollen originates from many species of the Poaceae family and release several proteins and glycoproteins,⁷ some of which are identified as allergens. These allergens are classified in 10 groups on the basis of their high intragroup cross-reactivity (see World Health Organization/International Union of Immunological Societies Allergen Nomenclature Sub-Committee at http://www.allergen.org/; Allergome at http://www.allergome.org/). *Phleum pratense* (timothy grass) is

From ^aZAUM-Center of Allergy & Environment, Helmholtz Center Munich/Technische Universität München, Member of the German Center for Lung Research (DZL), Munich; bthe Kühne Foundation, Christine Kühne-Center for Allergy Research and Education (CK-CARE), Munich; cthe Finnish Meteorological Institute, Erik Palmenin Aukio 1, Helsinki; ^dthe Laboratory of Allergology, Department of Clinical and Experimental Medicine, University of Parma, U.O. Medical Immunology, University Hospital of Parma; ethe EPAR Department, INSERM, UMR_S1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, France; ^fthe EPAR Department, Sorbonne Universités, UPMC Univ Paris 06, UMR_S 1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Paris; gICAAM-Institute of Mediterranean Crop and Environmental Sciences, University of Évora; hthe Center for Neuroscience and Cell Biology (CNC), University of Coimbra; ithe Medical University of Vienna, Department of Oto-Rhino-Laryngology, Research Unit Aerobiology and Pollen Information, Vienna; ^jthe Aerobiology Laboratory, Biology Department, Science and Arts Faculty, Uludag University, Gorukle-Bursa; ^kthe Department of Botany, Ecology and Plant Physiology, University of Córdoba, International Campus of Excellence on Agrifood (ceiA3), Córdoba; ¹the Laboratory of Aeropalynology, Faculty of Biology, Adam Mickiewicz University, Poznań; "the National Pollen and Aerobiology Research Unit, University of Worcester; "the University of Turku, CERUT, Aerobiology Unit, Turku; °Allergopharma GmbH & Co KG, Reinbek; Pthe Department of Environmental Research, Siauliai University, Siauliai; ^qRNSA (Réseau National de Surveillance Aérobiologique), Brussieu; and ^rthe Interdepartmental Centre of Bioclimatology, University of Florence.

Disclosure of potential conflict of interest: J. Buters has received or has grants pending from Foundation CK-CARE. S. Celenk's institution has received funding from the Scientific and Technological Research Council of Turkey (TUBITAK 109S032). B. Weber is employed by Allergopharma GmbH & Co. KG. The rest of the authors declare that they have no relevant conflicts of interest.

Received for publication January 22, 2014; revised January 7, 2015; accepted for publication January 12, 2015.

Corresponding author: Jeroen Buters, PhD, ZAUM–Center of Allergy & Environment, Helmholtz Center Munich/Technische Universität München, Biedersteinerstrasse 29, 80802 Munich, Germany. E-mail: buters@tum.de.

Abbreviations used					
HIALINE:	European Union Project Health Impacts of Airborne				
	Allergen Information Network				
10>PM>2.5:	Particulate matter larger than 2.5 µm and less than				
	10 μm in diameter				
PM>10:	Particulate matter larger than 10 µm in diameter				

a major source of grass pollen in temperate areas, and because of the cross-reactivity mentioned above, its extracts and single allergen components are widely used for the in vitro and in vivo diagnosis of grass pollen allergy and specific immunotherapy.⁸ Although sensitizations against Phl p 1, 2, 4, 5, 6, 7, 11, 12, and 13 are described, Phl p 1, 2, 5, and 6 are markers of grass pollen sensitization because they are present exclusively in grass pollen.^{8,9} Phl p 1, 4, and 5 show the highest rates of sensitization (>50%) in patients with grass pollen-induced allergy.^{10,11} However, recent findings have shown that the prevalence of sensitization and clinical importance of Phl p 4 might be overestimated.^{12,13} Other than for the clinical relevance, Phl p 5 is important for its high allergenic activity and good characterization.¹⁴ For these reasons, the variant Phl p 5.0102 (Phl p 5a) was adopted as reference material for the standardization of extracts for immunotherapy and skin prick tests, as indicated by the Development of Certified Reference Materials for Allergenic Products and Validation of Methods for Their Quantification (CREATE) project's findings.¹⁵ Different species of the Poaceae family express homologues of Phl p 5^{16,17} that show similar immunologic responses.¹

It is not possible to distinguish grass pollen grains from different grass species by using light microscopy, but the identity of allergens and the amount of allergen released from pollen varies between species.^{8,17} Natural variation of allergen from biological sources was reported for birch pollen,¹⁹ olive pollen,²⁰ horses,²¹ and cats.²² Allergen release also varies during the lifespan of pollen.²³

Monitoring allergens in ambient air enables the detection of non–pollen-bound allergens, as encountered during special weather conditions, such as thunderstorms.^{7,24} In addition, the determination of thresholds for allergic symptoms based on pollen counts has produced variable results^{25,26} and might be improved by adding airborne allergen monitoring.²⁷

In the framework of the European Union Project Health Impacts of Airborne Allergen Information Network (HIALINE; www.hialine.eu), we monitored daily amounts of grass pollen and their major group 5 allergens simultaneously across Europe. We showed that levels of Phl p 5 released from grass pollen was variable and emphasized the necessity to consider molecular aerobiology in addition to counting pollen for the assessment of exposure to airborne allergens, especially in clinical trials and epidemiologic studies.

METHODS

Pollen in ambient air

Airborne pollen concentrations were collected at 10 sites in Europe (Table I and see Table E1 and Fig E1 in this article's Online Repository at www.jacionline.org) by using Hirst-type volumetric spore traps, according to the requirements of the European Aeroallergen Network.²⁸ Quality of counts was controlled, as described previously.¹⁹

Allergen in ambient air

Air was simultaneously sampled at each site with ChemVol (Butraco, Son, The Netherlands) high-volume (800 L/min) cascade impactors equipped with dry polyurethane impacting substrates (that were prewashed with 0.1 mol/L ammonium bicarbonate and water) using the stages for particulate matter greater than 10 µm (PM>10) and particulate matter greater than 2.5 µm but less than 10 µm in diameter (10>PM>2.5), as described previously.²⁹ Allergen was extracted in 0.1 mol/L ammonium bicarbonate containing 0.1% BSA in an endover-end rotator at 100 rpm for 4 hours, lyophilized, and redissolved in one tenth of the original volume in PBS, according to standard procedures.^{23,30} Phl p 5 allergen was determined by means of ELISA with the mAbs 1D11 and BO1 (Allergopharma, Reinbek, Germany)³¹ calibrated against recombinant Phl p 5.0102.¹⁵ The assay recognizes group 5 allergens of Pooideae (Phl p 5.0102 and 5.0201); has no cross-reactivity with groups 1, 2, 3, 4, 6, and 13; and has a limit of detection of 1.0 ng of Phl p 5/mL, equaling about 2 pollen grains/ m^3 (1 mL of extract = 256 m³ of air). Response factors for the different group 5 allergens vary less than 2-fold (data not shown; Allergopharma, Reinbek, Germany). The assay recognizes C-terminal group 5 fragments, 32 and ELISA correlated with mass spectrometric quantification (data not shown, Allergopharma). Quality control of ELISA data was performed according to the rules developed in HIALINE.¹⁹ Values less than the detection limit were reported as zero.

Basophil degranulation assay

An immune cell response independent of ELISA was determined based on induction of mediator release by ambient air extracts from an FceRI-humanized rat basophil cell line (α -, β -, and γ -chain).³³ The cells were passively sensitized with human serum of a patient with grass pollen–induced symptoms, a positive SPT response, and a RAST score of greater than 3. The patient was sensitized to Phl p 1, 2/3, 4, 5b, 6, and 11 and Cyn d 1 but not Phl p 7 and 12. Levels of β -hexosaminidase as a proxy for histamine release was measured, as described previously.¹⁹ Each environmental sample was diluted sequentially, and only data from the linear part of the degranulation curve were reported.

Statistics

Pollen potency was determined several ways: (1) as the slope of the linear regression curve of pollen per cubic meter versus allergen per cubic meter with the intercept forced through zero and the strength of the relation expressed by the coefficient of correlation (r^2) ; (2) as the mode of the histogram of daily potencies; (3) as the median potency calculated for the whole data set and for subranges of daily pollen counts; and (4) as the 5th to 95th percentile range. The analysis excluded pollen counts of less than 10 grains/m³.^{19,34}

The station- and year-wise variability of potency was assessed from histograms for the individual sites. Differences were analyzed by using the Student *t* test, unless stated otherwise.³⁵ A *P* value of less than .05 was considered statistically significant. Values beyond 3 SDs of the mean were considered outliers.

The explanation factors for the potency variations included meteorological variables that were either monitored at the ChemVol locations or obtained from the nearest meteorological station.

Modeling

The System for Integrated Modelling of Atmospheric Composition (SILAM)^{36,37} was used to evaluate the flowering period and pollen atmospheric transport and also to compute the footprints of the observations, as described previously.¹⁹ The SILAM model was run with a time step of 15 minutes, and the footprints were calculated 60 hours backward in time for each daily observation at each site.

Daily observed potency was mapped to the origin of pollen through the footprints. The lowest values that make up 1% of the footprint integral were cut off to reflect the limited transport distance of grass pollen. The potency value was attributed to all areas covered by the footprint that flowered at the time of the footprint passage. The attributed potency values were then averaged over the season, finally resulting in potency maps that show the mean potency of pollen released from the area during a specific year.

2009-2011

		1999-2008*	2009		2010		2011		average ± SD (SD %)	
		Pollen index, Σ Grains/m³/γ	Pollen index, Σ Grains/m ³ /y	Potency, Phl p 5/grain (pg)	Pollen index, Σ Grains/m³/γ	Potency, Phl p 5/grain (pg)	Pollen index, Σ Grains/m ³ /y	Potency, Phl p 5/grain (pg)	Potency, Phl p 5/grain (pg)	
Austria	Vienna	3,024 ± 606†	NA	NA	2,604	3.546	2,411	2.941	3.244 ± 0.428 (13)	
Finland	Turku	740 ± 267	654	1.592	738	1.992	782	1.535	1.706 ± 0.249 (15)	
France	Varies [‡]	9,671 ± 3,808	10,420	2.405	7,130	2.403	6,708	2.937	2.582 ± 0.308 (12)	
Germany	Munich	$2,033 \pm 604$	2,041	3.042	1,821	2.127	2,184	2.133	2.434 ± 0.527 (22)	
Italy	Parma	7,521 ± 2,973	4,135	1.508	4,362	2.871	5,409	0.821	1.733 ± 1.043 (60)	
Poland	Poznan	4,697 ± 1,472§	6,078	2.703	7,898	3.398	4,482	1.996	2.699 ± 0.701 (26)	
Portugal	Evora	6,617 ± 8,836	5,725	2.144	17,113	1.499	22,815	1.545	1.729 ± 0.360 (21)	
Spain	Cordoba	5,609 ± 2,777	4,014	2.268	8,693	4.149	5,888	2.423	2.947 ± 1.044 (35)	
Turkey	Bursa	1,242 ± 490¶	1,881	NA	2,892	3.126	3,812	NA	3.126	
United Kingdom	Worcester	6,715 ± 2,223	4,885	5.906	5,141	3.570	3,531	4.178	4.551 ± 1.212 (27)	
Average potency	± SD			2.696 ± 1.395		2.868 ± 0.840		2.279 ± 0.989		
SD (%)				52		29		43		

TABLE I. Average yearly pollen count (pollen index) and allergen release per pollen for the stations in the different countries

Potency (release of allergen per pollen) was determined by the slope of the linear regression (see the Methods section). Means \pm SDs are presented. Exact location of the stations is given in Table E1.

NA, Not applicable.

*A complete preceding decade was not available for all stations, in which case the maximum period available was taken.

†Vienna 1999-2008: 2 sites: (A) 1999-2002 ATWIAK; (B) 2003-2008 ATWIEN. The trap was moved from ATWIAK (AKH–Allgemeines Krankenhaus) to ATWIEN (ZAMG– Zentralanstalt für Meteorologie und Geodynamik).

‡In 1999-2008, the station was in Lyon; in 2010, it was in Saint-Genis l'Argentière; and in 2011, it was in Brussieu.

§From 2005-2008.

||From 2001-2008.

¶From 1999-2000, 2003-2005, and 2008; no data from 2001-2002 and 2006-2007.

All simulations used the same configuration that included 8 vertical layers up to approximately 6 km above the ground. The horizontal grid cell size was 25 km. Input meteorological data were taken from the operational archive of the European Centre for Medium-Range Weather Forecast.

RESULTS

Pollen in ambient air

Airborne grass pollen is ubiquitous in Europe, but large variations in pollen counts exist between countries. A map of the stations is presented in Fig E1. The Finnish station in Turku (seaside) showed the lowest pollen counts, whereas the Portuguese station in Evora (rural site) recorded the highest counts (Table I). It should be noted that the years studied for each station were within the normal range of the preceding years for all stations (season timing and total pollen counts were within 1 SD from the long-term mean values, data not shown). The stations were distributed over Europe to obtain the most representative picture for the continent.

With 3 Hirst-type pollen traps operated simultaneously within 5 m of each other at about 9 m in height for a 3-week period during the grass pollen season (0-890 pollen grains/m³), we determined the variability between the traps to be about 5 pollen grains/m³ plus 20% (data not shown).

Allergen

In addition to variations in airborne pollen counts, pollen differed in allergen release per pollen (potency, Fig 1). Determined by means of linear regression (see the Methods section), the average group 5 allergen release per pollen was 2.3 pg of Phl p 5 per pollen ($r^2 = 0.497$, P < .001, n = 1629; see Fig E2 in this article's Online Repository at www.

jacionline.org). This number agrees well with the histogram mode (2-2.5 pg of Phl p 5 per pollen, see the histogram in Fig E3 in this article's Online Repository at www.jacionline. org). Median potency is somewhat greater, 2.6 pg of Phl p 5 per pollen, because of the asymmetric tails of the histogram. The pollen potencies for each station are given in Table I, and histograms are shown in Fig E4 in this article's Online Repository at www.jacionline.org. Potency distribution can have quite a wide spread that is different for different regions. Interannual variability of the histograms at the same station is smaller than spatial differences. Taking the extremes in pollen counts (Finland vs Portugal), allergen release per pollen was not different (not significant). Potency values over Europe were correlated with pollen counts (Fig 2).

We detected many days when pollen did not release any Phl p 5 at all. These pollen grains without group 5 allergens were especially noticeable at the beginning of the grass pollen seasons. This also explains why pollen potency distribution is not normally distributed (see Fig E3). Using the 5th to 95th percentile, pollen potency ranged between less than 1 and 9 pg of Phl p 5 per pollen (excluding low pollen counts [ie, <10 pollen grains/m³]).

Mediator release from basophils

For one station (Munich, Germany), we determined the daily mediator release by basophils passively sensitized to grass pollen allergens. In Munich no overlapping pollen season was measured, except for nonallergenic *Pinus* species (pine) and *Picea* species (spruce) pollen. The degranulation was standardized as if 1 m³ of air was delivered to the cells, which could result in calculated degranulation rates of greater than 100% (Fig 3). The serum used showed sensitization against several major and minor grass pollen

	2009	2010	2011
Austria		24 25 25 25 25 25 25 25 25 25 25	24 25 26 25 25 25 25 25 25 25 25 25 25
Finland	200 200 200 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 100 200 2	240 300 300 300 100 1	200 100 100 100 100 100 100 100
France	24 30 40 30 40 20 40 20 40 40 40 40 40 40 40 40 40 4		
Germany	250 200 110 110 100 100 100 100 10	24 20 10 10 10 10 10 10 10 10 10 1	250 200 101 101 101 101 101 101 10
Italy	24 30 40 40 40 40 40 40 40 40 40 4	24 20 4 4 5.4 13.4 13.4 13.4 13.4 13.5 15.5	20 10 10 10 10 10 10 10 10 10 1
Poland	24 30 5 30 5 5 5 5 5 5 5 5 5 5 5 5 5	24 20 4 17.5 31.5 146 286 127 28.5 9.6 10 17.5 31.5 146 28.6 127 28.5 9.6 10 17.5 31.5 146 28.5 127 28.5 9.6 10 17.5 31.5 146 28.5 127 28.5 10 17.5 31.5 127 38.5 10 17.5 31.5 10 17.5 31.	24 50 14 14 15 16 16 16 16 16 16 16 16 16 16
Portugal	2000 100 1000 1	200 100 100 100 100 100 100 100	200 100 100 100 100 100 100 100
Spain	500 500 500 500 500 500 500 500	24 20 40 40 40 40 40 40 40 40 40 4	90 90 90 90 90 90 90 90 90 90
Turkey		24 10 10 10 10 10 10 10 10 10 10	
UK	20 10 10 10 10 10 10 10 10 10 1	200 200 200 200 200 200 200 200	24 300 4 4 25. HAS. 30.5. IJA. 226. ILT. 257. RA. 228.

FIG 1. Pollen flight from all stations in 2009 until 2011 concomitant with pollen allergen release potency. *Gray vertical bars* represent pollen, and *purple lines* represent daily allergen released per pollen (daily pollen potency). *UK*, United Kingdom.



FIG 2. Experimental variability of grass pollen potency versus its estimated uncertainty. *Solid lines,* Potency uncertainty range calculated from assumed errors of pollen counts and ELISA. *Dashed lines,* Observed percentiles; *thick dots* indicate the centers of bins used for percentiles estimation. The observed potency variability was much larger than can be explained by experimental error. Pollen counts of less than 10 pollen grains/m³ were excluded.

allergens (Phl p 1, 2/3, 4, 5b, 6, and 11). Although these specific IgEs to several major or minor allergens could have mediated basophil degranulation, mediator release from basophilic granulocytes correlated well with Phl p 5 levels per cubic meter ($r^2 = 0.80$, P < .001) and with pollen levels per cubic meter ($r^2 = 0.61$, P < .001). Two outliers (probably because of the large dilution needed to achieve the linear part of the degranulation curve) were identified and indicated in Fig 3.

Potency maps

Potency maps compiled for each year (Fig 4, left column) showed a substantial variability over Europe. Sampling stations were located between 2 and 22 m (see Table E1). Fewer grass pollens were sampled at higher monitor locations because of the substantial sedimentation velocity of grass pollen. However, numeric simulations showed that the separation of different sizes along the height is negligible (<20% in the most extreme cases) at the used heights (data not shown). Thus the differences in sampling height in our study do not explain the differences in pollen potency.

Several regions, such as France and Germany, appeared to have quite high potency of the pollen released, reaching values greater than 5 pg of Phl p 5 per pollen. The lowest potency in all years was registered in Eastern Europe in 2011 at less than 2 pg of Phl p 5 per pollen. However, most regions appeared to vary from year to year.

A crude estimate of reliability of these maps can be obtained from the SD of potency (Fig 4, right column). The high potency estimated in France is usually accompanied by a large spread that is similar but less than the values themselves. The stripes visible on the maps are individual footprints corresponding to substantially different observed potencies. Their overlap results in high uncertainty of the final pattern. In other regions, however, the uncertainty was lower and independent of the actual potency value.

Relation of potency and pollen counts

The average pollen potency stayed almost constant throughout the observed pollen count range (Fig 2, thick black



FIG 3. Degranulation of FccRI-humanized rat basophils by extracts of ambient air. **A** and **B**, Degranulation caused by allergen of PM>10 (Fig 3, A) or degranulation caused by pollen (Fig 3, B). Two extremes indicated in the graph were excluded in both calculations of correlation (see the Methods section).

dashed line denotes its median). However, slight decreasing trends are noticeable for low and high counts. To the contrary, the spread clearly increases toward low counts. One can estimate what fraction of this variability is due to observational errors and what is to be attributed to natural potency variability. As stated above, the experimental error for pollen counts is about 5 pollen grains/m³ plus 20% of the observed value, and the error of ELISA is approximately 20%. Because potency distribution is non-Gaussian, robust estimates of the 1 and 2 SD ranges will be the differences between the potency values at 16% to 84% and 2% to 98%, respectively. As one can see, pollen varied more in potency than can be explained by the experimental errors.

Humidity

Allergen was collected in 2 fractions of ambient air. Within all of Europe, more than $89\% \pm 11\%$ of the allergen was collected as PM>10. The geometric diameter of grass pollen is 20 to 40 μ m,³⁸ exemplifying that allergen in ambient air coincides with pollen. However, on some days, allergen in 10>PM>2.5,



FIG 4. Map of Europe showing the potency of pollen to release allergen in 2009 (**A**), 2010 (**B**), and 2011 (**C**), depending on the pollen origin (source). Source attribution is needed because ripening of allergen occurs at the source. Concomitantly, the SD in the corresponding years is given. Pollen potency is not fixed to certain regions and changes between years.



FIG 5. Phl p 5 as 10>PM>2.5 versus total Phl p 5 in ambient air (total, ratio in percentages) for all stations for all years, depending on local relative humidity, showing allergen in the non–pollen-containing fraction of ambient air. Zero group 5 allergen values and values of less than 10 pollen grains/m³ were excluded. Means \pm SDs are presented (*P* < .001, n = 1252).

where normally few pollen is detected, reached 20% of total airborne allergen. The allergen content of this fraction of air can contain smaller particles, which correlated with humidity (P < .001, Fig 5).

DISCUSSION

Pollen is the major outdoor cause for allergic rhinitis, and more than 350 pollen traps over Europe monitor pollen flight daily.³⁹ The immune system reacts with symptoms to allergens,⁴⁰ which are carried and released by pollen. The natural variability across Europe of how much allergen is carried by grass pollen was unknown.

This study shows that, on average, grass pollen in Europe released about 2.0 to 2.5 pg of group 5 allergens, the major grass pollen allergen, per pollen grain. This average value has a large variation because the amount of group 5 allergen released by the same amount of pollen ranged from less than 1 pg (detection limit) to 9 pg per pollen (5th to 95th percentile) and varied 7-fold between countries and years (P < .01).

Although pollen counts are indispensable for agriculture, phenology, climate change, and health, airborne allergen adds a new dimension to understanding allergic rhinitis and asthma. Indeed, depending on humidity, up to 20% of the allergen could be detected in the fraction of air containing smaller particles than pollen that could penetrate deeper into the lungs.

Pollen in ambient air correlated with allergic symptoms,^{26,40} also for grass pollen.⁴¹ However, the correlation is not clear because symptoms lag behind exposure and later in the season higher pollen counts do not evoke more symptoms. The human immune system reacts to allergens with symptoms, and allergen

release of pollen is not constant (Fig 1 and Table I). In addition, more events with high-potency pollen were observed when pollen counts were low (Fig 2). This could explain why even low levels of 2 to 9 pollen grains/m³ can result in allergic symptoms.⁴² Also, at high humidity, more allergen was detected in the fraction of air containing smaller particles that can penetrate deeper into the lungs⁴³ (Fig 5; ie, depending on humidity, a part of the allergen could penetrate deeper into the airways, approximately double at 100% relative humidity [rainy days] vs dry days with 30% relative humidity; P < .001). As a consequence, pollen concentration predicted allergen exposure but only with an r^2 correlation of 0.497 (P < .001, see Fig E2).

Allergen in ambient air

For birch pollen, a 10-fold natural variation in allergen release per pollen was published; for olive pollen, this was 12-fold.^{19,20} For the sake of comparing the allergen release variability for grass with those results, we also followed the approach of those studies and compared the mean potency of the top 10% of values with the lowest 10%. For grass, this additionally required excluding pollen without group 5 allergens and pollen with potencies of greater than 20 pg Phl p 5 per pollen. Also excluded were days with pollen counts of less than 10 pollen grains/m³. Then the variability in potency for grass pollen was estimated to be 17-fold. With all the ambiguity of such a comparison, the range of variations among the considered taxa appeared to be the largest for grass.

A weakness of this study is that no other grass pollen allergens, particularly group 1 allergens, were determined. Grass pollen exposure involves a succession of pollen from different grass species.⁴⁴ Grass pollen of different species release similar, crossreacting group 5 allergens that are not identical (www.allergome. org, accessed August 2013). Thus Dactylis glomerata pollen releases Dac g 5, whereas P pratense releases Phl p 5. The response factor of each Phl p 5 analogue is different,¹⁸ as reported for other allergens, such as Phl p 1.¹⁷ We calibrated all samples against Phl p 5.0102, analogue to calibrating protein concentration against BSA,⁴⁵ because Phl p 5.0102 is the pharmacopeia standard for Phleum species preparations used in specific immunotherapy.³¹ For group 5 allergens, the difference in response factor between the early flowering major species Dglomerata and the late blooming species P pratense is low, maximally about 2-fold, with Phl p 5 having the lower response factor. Differences in response factor are not sufficient to explain the observed difference in pollen potency (range, <1-9 pg of Phl p 5 per pollen [5th to 95th percentile]).¹⁷

The flowering succession of grass species includes, among other minor species, *Alopecurus pratensis*, *Poa pratensis*, *Lolium perenne*, *D glomerata*, and *P pratense*.⁴⁴ *A pratensis* (beginning of the season) and *Phragmites* species (end of season) do not release Phl p 5 (and data not shown),^{8,18} and indeed, at the beginning of the season, several stations detected pollen without group 5 allergens (Fig 1 and see Figs E3 and E4). For instance, the lower potency of pollen in Munich at the beginning of the grass pollen season in 2011 (Fig 1) was not due to a lower response factor because *D glomerata* flowers before *P pratense* and has a higher response factor to Phl p 5.0102 but must be due to a lower release of total group 5 allergens per pollen.

Analogous to having pollen without group 5 allergens at the beginning of the grass pollen season, at the end of the grass pollen

season in Finland, where *Phragmites* species (common reed) are more abundant, also pollen with no Phl p 5 is detected. In other countries the contribution to pollen counts by this species is limited.^{18,46}

Mediator release

In the basophil degranulation bioassay β -hexosaminidase as a proxy for histamine release by the particle equivalent of 1 m^3 of ambient air correlated well with allergen levels per cubic meter $(r^2 = 0.80, P < .001)$ and pollen levels per cubic meter $(r^2 = 0.61, P < .001, Fig 3)$. Pollen releases more allergymodulating factors than allergens.⁴⁷ The degranulation assay also agrees with our hypothesis that pollen releases different amounts of Phl p 5, as reported previously for other pollen species.^{19,20} However, the basophil assay is not a substitute for allergy symptoms. It should also be taken into account that the reliability of the group 5 ELISA as a predictor of biological activity depends on the variability of the group 1/group 5 ratio. This highly significant correlation presumably indicates that the group 1/group 5 ratio in our air samples, which were all from the Munich area, is relatively constant and that the contribution of other allergens is small. We cannot exclude that pollen collected in Munich might have a different and/or more stable group 1/group 5 ratio than pollen collected in other parts of Europe.

Potency variations in time and space

Pollen grains varied in the amount of allergen they release. However, most data stem from single samples and might not be representative because of interday, intercountry, and interyear variability of the natural sources.^{17,18,48} In addition, allergen levels throughout the literature were quantified with different ELISAs, yielding variable results. Therefore we used the Development of Certified Reference Materials for Allergenic Products and Validation of Methods for Their Quantification proposed methods and reference materials,^{15,49} eliminating this variability.

Considering the potency maps and their SDs, one should keep in mind the peculiarity of grass pollen: in different regions and at different times, it is released by different species, which are likely to have different potencies. This is corroborated by substantial variability between the monthly potency maps for each season (data not shown). In Fig 4 this temporal variability contributes to the overall seasonal SD (Fig 4, right column).

At the source, variability in pollen potency can be the result of 2 ripening processes occurring simultaneously. First, pollens increase their allergen content rapidly on ripening inside the anthers just before pollination.²³ Second, when anthers have ripened sufficiently, they react to actual weather conditions, releasing pollen.⁵⁰ These parallel ripening processes mean that the amount of allergen in the released pollen depends on the actual weather before and during the flowering season, resulting in weather-dependent variations in potency at the monitoring site.

Pollens varied more in their potency than can be explained by experimental error (Fig 2). Potency has an uncertainty stemming from both uncertainty in pollen count and uncertainty in Phl p 5 determinations. For pollen counts, the inaccuracy is poorly known,⁵¹ and we estimated it as 5 pollen grains/m³ plus 20% of the observed value, which is similar to the inaccuracy for birch pollen counts.¹⁹ For ELISA in this study, the inaccuracy was 20.4%, which is similar, as generally reported for ELISAs.^{19,52} Using both uncertainties, we showed that the observed potency variability was several-fold higher than what could be explained by the experimental error (Fig 2). Because we determined the amount of allergen in both fractions, PM>10 and 10>PM>2.5 of ambient air, 2 explanation are possible. Either pollens with low pollen counts release more allergen or more allergen was already present as 10>PM>2.5, which could be free allergen.

Humidity

At higher relative humidity of ambient air during pollination, more allergen was detected as 10>PM>2.5 (Fig 5). Pollen grains were detected in this fraction of air, despite their large size, because of incomplete separation by means of cascade impaction.^{29,53} At higher humidity, either more pollen was collected as 10>PM>2.5 because of pollen grains changing their aerodynamic diameters or more allergen is present in the form of smaller particles. In contrast to other observations, which described extreme weather conditions, such as thunderstorms, when these effects appeared,^{24,54,55} we could show that by less extreme weather conditions, such as higher humidity, more allergen, either as changed pollen or as smaller particles, was available in ambient air that could penetrate deeper into the lung, evoking more severe symptoms. As grass pollen swells and increases its aerodynamic diameter on exposure to higher humidity (A. LeMoal, unpublished observation), we think the presence of more small Phl p 5-containing particles at higher humidity is the most likely explanation. This hypothesis was also suggested by others⁴¹ and even more closely related to the results of Schäppi et al⁴⁸ after light rainfall condition.

The results indicate that the pollen potency is not constant and, in addition, could be explained by a free fraction of allergen in ambient air that could depend on humidity and is more easily detected when less pollen is in the air (Figs 2 and 5).

Conclusion

Pollens across Europe varied in their natural capacity to release group 5 allergens. This is in addition to the natural variation in pollen counts. In a biological test system mediator release correlated better with ambient allergen concentrations than with pollen counts. No clear geographic pattern of pollen potency was detected. Although pollen is an excellent proxy for exposure, more can be learned from the actual monitoring of allergens in ambient air. Indeed, at higher humidity, more allergen seems to be present on smaller particles that could penetrate deeper into the airways.

The technical assistance of Danijel Kupresanin, Ingrid Weichenmeier, Christine Huber, Amelie Le Moal-Le Berre, Elsa Caeiro, Raquel Ferro, Manuela Ugolotti, Małgorzata Nowak, Agata Szymańska, Łukasz Kostecki, and Satu Saaranen is greatly appreciated. We thank Robert Gebauer for the construction of the database and Dr S. Vieths for donating the humanized basophils. The excellent management assistance of Annina Sorgner was pivotal for the project.

Key messages

- The same amount of pollen can release variable amounts of allergen ranging from less than 1 to 9 pg of Phl p 5 per pollen (5th to 95th percentile).
- This difference varies per location in Europe and during the pollen season.
- At higher humidity, the amount of group 5 allergen not associated with pollen increases.

REFERENCES

- Burbach GJ, Heinzerling LM, Edenharter G, Bachert C, Bindslev-Jensen C, Bonini S, et al. GA(2)LEN skin test study II: clinical relevance of inhalant allergen sensitizations in Europe. Allergy 2009;64:1507-15.
- Sunyer J, Jarvis D, Pekkanen J, Chinn S, Janson C, Leynaert B, et al. Geographic variations in the effect of atopy on asthma in the European Community Respiratory Health Study. J Allergy Clin Immunol 2004;114:1033-9.
- Jarvis D, Luczynska C, Chinn S, Potts J, Sunyer J, Janson C, et al. Change in prevalence of IgE sensitization and mean total IgE with age and cohort. J Allergy Clin Immunol 2005;116:675-82.
- Scala E, Alessandri C, Bernardi ML, Ferrara R, Palazzo P, Pomponi D, et al. Cross-sectional survey on immunoglobulin E reactivity in 23,077 subjects using an allergenic molecule-based microarray detection system. Clin Exp Allergy 2010;40:911-21.
- Haftenberger M, Laussmann D, Ellert U, Kalcklosch M, Langen U, Schlaud M, et al. [Prevalence of sensitisation to aeraoallergens and food allergens: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2013;56: 687-97.
- Canova C, Heinrich J, Anto JM, Leynaert B, Smith M, Kuenzli N, et al. The influence of sensitisation to pollens and moulds on seasonal variations in asthma attacks. Eur Respir J 2013;42:935-45.
- Abou Chakra OR, Sutra JP, Demey Thomas E, Vinh J, Lacroix G, Poncet P, et al. Proteomic analysis of major and minor allergens from isolated pollen cytoplasmic granules. J Proteome Res 2012;11:1208-16.
- Gangl K, Niederberger V, Valenta R. Multiple grass mixes as opposed to single grasses for allergen immunotherapy in allergic rhinitis. Clin Exp Allergy 2013; 43:1202-16.
- Andersson K, Lidholm J. Characteristics and immunobiology of grass pollen allergens. Int Arch Allergy Immunol 2003;130:87-107.
- Tripodi S, Frediani T, Lucarelli S, Macri F, Pingitore G, Di Rienzo Businco A, et al. Molecular profiles of IgE to *Phleum pratense* in children with grass pollen allergy: implications for specific immunotherapy. J Allergy Clin Immunol 2012; 129:834-9.
- Moverare R, Kosunen TU, Haahtela T. Change in the pattern of IgE reactivity to timothy grass and birch pollen allergens over a 20-year period. J Investig Allergol Clin Immunol 2006;16:274-8.
- Westritschnig K, Horak F, Swoboda I, Balic N, Spitzauer S, Kundi M, et al. Different allergenic activity of grass pollen allergens revealed by skin testing. Eur J Clin Invest 2008;38:260-7.
- Zafred D, Nandy A, Pump L, Kahlert H, Keller W. Crystal structure and immunologic characterization of the major grass pollen allergen Phl p 4. J Allergy Clin Immunol 2013;132:696-703.
- 14. Flicker S, Vrtala S, Steinberger P, Vangelista L, Bufe A, Petersen A, et al. A human monoclonal IgE antibody defines a highly allergenic fragment of the major timothy grass pollen allergen, Phl p 5: molecular, immunological, and structural characterization of the epitope-containing domain. J Immunol 2000; 165:3849-59.
- Chapman MD, Ferreira F, Villalba M, Cromwell O, Bryan D, Becker WM, et al. The European Union CREATE project: a model for international standardization of allergy diagnostics and vaccines. J Allergy Clin Immunol 2008;122:882-9.
- Bullimore A, Swan N, Alawode W, Skinner M. A proteomic style approach to characterize a grass mix product reveals potential immunotherapeutic benefit. World Allergy Organ J 2011;4:140-6.
- Duffort O, Quintana J, Ipsen H, Barber D, Polo F. Antigenic similarity among group 1 allergens from grasses and quantitation ELISA using monoclonal antibodies to Phl p 1. Int Arch Allergy Immunol 2008;145:283-90.
- Ramirez J, Obispo TM, Duffort D, Carpizo JA, Chamorro MJ, Barber D, et al. Group 5 determination in Pooideae grass pollen extracts by monoclonal

antibody-based ELISA. Correlation with biologic activity. Allergy 1997;52: 806-13.

- Buters JTM, Thibaudon M, Smith M, Kennedy R, Rantio-Lehtimaaki A, Albertini R, et al. Release of Bet v 1 from birch pollen from 5 European countries. Results from the HIALINE study. Atmos Environ 2012;55:496-505.
- 20. Galan C, Antunes C, Brandao R, Torres C, Garcia-Mozo H, Caeiro E, et al. Airborne olive pollen counts are not representative of exposure to the major olive allergen Ole e 1. Allergy 2013;68:809-12.
- Mitlehner W. Allergy against horses. Are curly horses an alternative for horse-allergic riders? Allergo J 2013;22:244-51.
- 22. Bienboire-Frosini C, Cozzi A, Lafont-Lecuelle C, Vervloet D, Ronin C, Pageat P. Immunological differences in the global release of the major cat allergen Fel d 1 are influenced by sex and behaviour. Vet J 2012;193:162-7.
- 23. Buters JT, Weichenmeier I, Ochs S, Pusch G, Kreyling W, Boere AJ, et al. The allergen Bet v 1 in fractions of ambient air deviates from birch pollen counts. Allergy 2010;65:850-8.
- D'Amato G, Cecchi L, Liccardi G. Thunderstorm-related asthma: not only grass pollen and spores. J Allergy Clin Immunol 2008;121:537-8.
- de Weger L, Dahl A, Bergmann K, Belmonte J, Rantio-Lehtimaaki A, Thibaudon M, et al. Impact of pollen. In: Bergman CK, Sofiev M, editors. Allergenic pollen. New York: Springer; 2012.
- 26. Caillaud DM, Martin S, Segala C, Besancenot JP, Clot B, Thibaudon M. Nonlinear short-term effects of airborne Poaceae levels on hay fever symptoms. J Allergy Clin Immunol 2012;130:812-4.
- Cecchi L. From pollen count to pollen potency: the molecular era of aerobiology. Eur Respir J 2013;42:898-900.
- 28. European Aerosol Society. Minimum requirements to manage aerobiological monitoring stations included in a national network involved in the EAN. International Aerobiology Newsletter 2011;72:1.
- Demokritou P, Gupta T, Ferguson S, Koutrakis P. Development and laboratory characterization of a prototype coarse particle concentrator for inhalation toxicological studies. J Aerosol Sci 2002;33:1111-23.
- **30.** Buters JT, Kasche A, Weichenmeier I, Schober W, Klaus S, Traidl-Hoffmann C, et al. Year-to-year variation in release of Bet v 1 allergen from birch pollen: evidence for geographical differences between West and South Germany. Int Arch Allergy Immunol 2008;145:122-30.
- 31. Vieths S, Barber D, Chapman M, Costanzo A, Daas A, Fiebig H, et al. Establishment of recombinant major allergens Bet v 1 and Phl p 5a as Ph. Eur. reference standards and validation of ELISA methods for their measurement. Results from feasibility studies. Pharmeur Bio Sci Notes 2012;2012:118-34.
- **32.** van Oort E, Dieker MC, de Heer PG, Peltre G, Aalberse RC, van Ree R. The major grass pollen group 5 allergen from *Dactylis glomerata* and its C-terminal split product both behave as dimers: implications for allergen standardization. Int Arch Allergy Immunol 2005;136:113-22.
- 33. Vogel L, Luttkopf D, Hatahet L, Haustein D, Vieths S. Development of a functional in vitro assay as a novel tool for the standardization of allergen extracts in the human system. Allergy 2005;60:1021-8.
- Lorenz RJ. Biometrie. Grundbegriffe der Biometrie. 2nd ed. Stuttgart: Gustav Fischer Verlag; 1989.
- 35. Livingston EH. Who was Student and why do we care so much about his t-test? J Surg Res 2004;118:58-65.
- 36. Prank M, Chapman DS, Bullock JM, Belmonte Soler J, Berger U, Dahl A, et al. An operational model for forecasting ragweed pollen release and dispersion in Europe. Agric For Meteorol 2013;182-183:43-53.

- 37. Sofiev M, Siljamo P, Ranta H, Linkosalo T, Jaeger S, Rasmussen A, et al. A numerical model of birch pollen emission and dispersion in the atmosphere. Description of the emission module. Int J Biometeorol 2013;57:45-58.
- Brown HM, Irving KR. The size and weight of common allergenic pollens. An investigation of their number per microgram and size distribution. Acta Allergol 1973;28:132-7.
- 39. Berger U, Karatzas K, Jaeger S, Voukantsis D, Sofiev M, Brandt O, et al. Personalized pollen-related symptom-forecast information services for allergic rhinitis patients in Europe. Allergy 2013;68:963-5.
- Jantunen J, Saarinen K, Rantio-Lehtimaaki A. Allergy symptoms in relation to alder and birch pollen concentrations in Finland. Aerobiologia 2012;28:169-76.
- 41. Brito F, Mur Gimeno P, Carnes J, Fernandez-Caldas E, Lara P, Alonso A, et al. Grass pollen, aeroallergens, and clinical symptoms in Ciudad Real, Spain. J Investig Allergol Clin Immunol 2010;20:295-302.
- DellaValle CT, Triche EW, Leaderer BP, Bell ML. Effects of ambient pollen concentrations on frequency and severity of asthma symptoms among asthmatic children. Epidemiology 2012;23:55-63.
- 43. International Commission on Radiological Protection. Human respiratory tract model for radiological protection, ICRP publication 66. New York: Pergamon Press, Elsevier Science; 1994.
- 44. Frenguelli G, Passalacqua G, Bonini S, Fiocchi A, Incorvaia C, Marcucci F, et al. Bridging allergologic and botanical knowledge in seasonal allergy: a role for phenology. Ann Allergy Asthma Immunol 2010;105:223-7.
- 45. Peterson GL. A simplification of the protein assay method of Lowry et al. which is more generally applicable. Anal Biochem 1977;83:346-56.
- 46. Leon-Ruiz E, Alcazar P, Dominguez-Vilches E, Galan C. Study of Poaceae phenology in a Mediterranean climate. Which species contribute most to airborne pollen counts? Aerobiologia 2011;27:37-50.
- Gilles S, Fekete A, Zhang X, Beck I, Blume C, Ring J, et al. Pollen metabolome analysis reveals adenosine as a major regulator of dendritic cell-primed T(H) cell responses. J Allergy Clin Immunol 2011;127:454-61.
- 48. Schäppi GF, Taylor PE, Pain MC, Cameron PA, Dent AW, Staff IA, et al. Concentrations of major grass group 5 allergens in pollen grains and atmospheric particles: implications for hay fever and allergic asthma sufferers sensitized to grass pollen allergens. Clin Exp Allergy 1999;29:633-41.
- 49. van Ree R, Chapman MD, Ferreira F, Vieths S, Bryan D, Cromwell O, et al. The CREATE project: development of certified reference materials for allergenic products and validation of methods for their quantification. Allergy 2008;63:310-26.
- Taylor PE, Flagan RC, Valenta R, Glovsky MM. Release of allergens as respirable aerosols: a link between grass pollen and asthma. J Allergy Clin Immunol 2002;109:51-6.
- Bergmann KC, Simoleit A, Wagener S, Mücke HG, Werchan M, Zberbier T. Verteilung von Pollen und Feinstaub in einem städtischen Ballungsraum am Beispiel der großstadt Berlin. Allergo J 2013;22:471-5.
- King EM, Filep S, Smith B, Platts-Mills T, Hamilton RG, Schmechel D, et al. A multi-center ring trial of allergen analysis using fluorescent multiplex array technology. J Immunol Methods 2013;387:89-95.
- Pöttering H, Lenarcic J. Directive 2008/50/EC of the European Parliament and of the council. Official Journal of the European Union 2008;152:1-44.
- Taylor PE, Jonsson H. Thunderstorm asthma. Curr Allergy Asthma Rep 2004;4: 409-13.
- Venables KM, Allitt U, Collier CG, Emberlin J, Greig JB, Hardaker PJ, et al. Thunderstorm-related asthma—the epidemic of 24/25 June 1994. Clin Exp Allergy 1997;27:725-36.



FIG E1. Map of station locations used in HIALINE.

Potency pollen across Europe with lin. regression



FIG E2. Pollen flight from all stations in 2009 until 2011 concomitant with airborne allergen concentration. Pollen values of less than 10 pollen grains/m³ were deleted for this correlation. The slope of the Pearson linear regression curve is given and represents allergen release per pollen (potency, 2.3 pg PhI p 5/pollen, r²=0.495, *P* < .001, n = 1629).



FIG E3. Histogram of grass pollen potency across Europe. The high frequency of low-potency pollen was dominated by pollen releasing no PhI p 5.



 $\ensuremath{\text{FIG}}$ E4. Histograms of grass pollen potency for each observation site and year.



FIG E4. (Continued).

TABLE E1. Station descriptions

		Location	Above ground (m)	Above sea level (m)
Austria	Vienna	48°14′56.0 ^{′′} North, 16°21′22.0 ^{′′} East	9.0	209
Finland	Turku	60°27′18.34 ^{''} North, 22°17′07.49 ^{'''} East	15	40
France	Lyon 2009-2010*	45°42′40.3 ^{''} North, 4°29′34.6 ^{''} East	2	478
	Lyon 2010†	45°45′09.5″ North, 4°32′17.9″ East	2	448
Germany	Munich	48°09′52.3 ^{''} North, 11°35′35.4 ^{''} East	1.8	520
Italy	Parma	44°48′16.2 ^{''} North, 10°18′56.9 ^{''} East	18	52
Poland	Poznan	52°28′01.8 ^{''} North, 16°55′27.3 ^{''} East	18	93
Portugal	Evora	38°34′07.2 ^{′′} North, 7°54′68.9 ^{′′} West	12	300
Spain	Cordoba	37°53′0.5″ North, 4°46′45.0″ West	22	123
Turkey	Bursa	40°13′21.2 ^{''} North, 28°51′48.2 ^{''} East	9	169
United Kingdom	Worcester	52°19'69.0 ^{''} North, 2°24'20.0 ^{''} West	10	43

ChemVol and Hirst-type pollen trap were located at the same location at heights within 5 m of each other. *Saint-Genis l'Argentiere. †Brussieu (8 km apart).